Contracting organization	Mascot International A/S, Denmark		
Project team	Corporate Responsibility Department, Mascot International A/S		
Review of Mascot's Life-Cycle Assessment (LCA) methodology and product LCA	Quantis Sàrl, Switzerland		
Method validity date	December 2023 Methodology is valid for 5 years		
Method	ISO 14040:2006 + A1:2020 / ISO 14044:2006 + A1:2018 + A2:2020. Product Environmental Footprint Category Rules (PEFCR) for Apparel and Footwear is followed when possible.		
Description of system boundaries	Cradle to grave		
LCIA method	EF 3.1 (adapted)		
Data collection	Primary data – main source. Generic data from ecoinvent v.3.10 APOS database Reference year is 2023		
LCA software used	SimaPro v.9.6.0.1		
Data quality	Method for data quality rating (DQR) developed in alignment with the PEF requirements.		
Data quality declaration	High (rated as described in PEFCR for Apparel and Footwear).		
Limitations	Style studies are based on reference sizes as defined in PEFCR for apparel and footwear. Current model is also based on reference colours. For other sizes and colours, the reader is encouraged to bear this in mind.		
LCA methodology summary report	Contact <u>responsibility@mascot.dk</u> if you are interested in the report.		

Main fabric: 88% PES / 12% EOL

# LIFE CYCLE ASSESSMENT FACTSHEET

March 2025 version 2.1

## **TARGET GROUP**

The 18008 is part of a collection designed for a broad target group in different work situations within trade, construction, manufacturing, industry and businesses with laundry agreements.

### LONG-LASTING DURABILITY

By analysing fabric performance requirements and collecting data on customer experience, the LCA is verified by Quantis for an estimated duration of service of use in hard working situations and with industrial wash every week.

#### **CRADLE-TO-GRAVE**

Cradle-to-grave is a scoping of the LCA that calculates the entire lifecycle of a product from Extraction of Raw materials to the Use & Wash and End-of-Life stages Cradle-to-grave results are presented per use according to PEF Category Rules for Apparel and Footwear.


### **METHODOLOGY**

MASCOT LCAs is mainly based on primary data from own factories and suppliers. MASCOT LCAs are calculated according to ISO14040/44. The method is verified by Quantis and applies to all colours.




Cradle to Grave: 0,055kg CO₂ per use Based on an ISO compliant methodology verified by Quantis

# **PROCESS CHAIN**







Main fabric: 88% PES / 12% EOL

# THE 16 IMPACT FACTORS

Impact category	Damage assessment	Impact to-gate	Impact to-grave
	unit	per garment	per use
Acidification	mol H⁺ eq	0,0481	0,000226
Climate change	kg CO <sub>2</sub> eq	8,1	0,055
Climate change - Biogenic	kg CO₂ eq	0,0125	0,00056
Climate change - Fossil	kg CO₂ eq	8,08	0,0534
Climate change - Land use and LU change	kg CO₂ eq	0,00735	0,000994
Ecotoxicity, freshwater	CTUe	30	0,455
Ecotoxicity, freshwater - part 2	CTUe	27,7	0,136
Ecotoxicity, freshwater - inorganics	CTUe	49,8	0,462
Ecotoxicity, freshwater - organics part 1	CTUe	5,62	0,116
Ecotoxicity, freshwater - organics part 2	CTUe	2,29	0,0132
Particulate matter	disease inc.	0,00000045	0,00000000212
Eutrophication, marine	kg N eq	0,00887	0,0000672
Eutrophication, freshwater	kg P eq	0,000384	0,00000576
Eutrophication, terrestrial	mol N eq	0,0959	0,000519
Human toxicity, cancer	CTUh	0,0000000254	0,00000000173
Human toxicity, cancer - inorganics	CTUh	0,0000000154	0,00000000000675
Human toxicity, cancer - organics	CTUh	0,0000000238	0,00000000166
Human toxicity, non-cancer	CTUh	0,00000136	0,000000000627
Human toxicity, non-cancer - inorganics	CTUh	0,000000125	0,00000000578
Human toxicity, non-cancer - organics	CTUh	0,000000115	0,000000000488
Ionising radiation	kBq U <sup>-235</sup> eq	0,567	0,00214
Land use	Pt	32,5	0,254
Ozone depletion	kg CFC11 eq	0,0000103	0,0000000248
Photochemical ozone formation	kg NMVOC eq	0,0373	0,000197
Resource use, fossils	MJ	128	0,799
Resource use, minerals and metals	kg Sb eq	0,000154	0,000000481
Water use	m³ depriv.	13,8	0,0412

