LCA Information

Style nr. 18053-316

	and the second		
Contracting organization	Mascot International A/S, Denmark		
Project team	Corporate Responsibility Department, Mascot International A/S		
Review of Mascot's Life-Cycle Assessment (LCA) methodology and product LCA	Quantis Sàrl, Switzerland		
Method validity date	December 2023 Methodology is valid for 5 years		
Method	ISO 14040:2006 + A1:2020 / ISO 14044:2006 + A1:2018 + A2:2020. Product Environmental Footprint Category Rules (PEFCR) for Apparel and Footwear is followed when possible.		
Description of system boundaries	Cradle to grave		
LCIA method	EF 3.1 (adapted)		
Data collection	Primary data – main source. Generic data from ecoinvent v.3.10 APOS database Reference year is 2023		
LCA software used	SimaPro v.9.6.0.1		
Data quality	Method for data quality rating (DQR) developed in alignment with the PEF requirements.		
Data quality declaration	High (rated as described in PEFCR for Apparel and Footwear).		
Limitations	Style studies are based on reference sizes as defined in PEFCR for apparel and footwear. Current model is also based on reference colours. For other sizes and colours, the reader is encouraged to bear this in mind.		
LCA methodology summary report	Contact <u>responsibility@mascot.dk</u> if you are interested in the report.		

LIFE CYCLE ASSESSMENT FACTSHEET

March 2025 version 2.1

TARGET GROUP

The 18053 is part of a collection designed for a broad target group in different work situations within trade, construction, manufacturing, industry and businesses with laundry agreements.

LONG-LASTING DURABILITY

By analysing fabric performance requirements and collecting data on customer experience, the LCA is verified by Quantis for an estimated duration of service of use in hard working situations and with industrial wash every week.

CRADLE-TO-GRAVE


Cradle-to-grave is a scoping of the LCA that calculates the entire lifecycle of a product from Extraction of Raw materials to the Use & Wash and End-of-Life stages. Cradle-to-grave results are presented per use according to PEF Category Rules for Apparel and Footwear.

METHODOLOGY

MASCOT LCAs is mainly based on primary data from own factories and suppliers. MASCOT LCAs are calculated according to ISO14040/44. The method is verified by Quantis and applies to all colours.



PROCESS CHAIN





WORKWEAR

THE 16 IMPACT FACTORS

Impact category	Damage assessment	Impact to-gate	Impact to-grave
	unit	per garment	per use
Acidification	mol H⁺ eq	0,024	0,000139
Climate change	kg CO ₂ eq	4,46	0,037
Climate change - Biogenic	kg CO ₂ eq	0,00949	0,000502
Climate change - Fossil	kg CO ₂ eq	4,45	0,0358
Climate change - Land use and LU change	kg CO ₂ eq	0,00556	0,000736
Ecotoxicity, freshwater	CTUe	23,7	0,342
Ecotoxicity, freshwater - part 2	CTUe	17,3	0,0928
Ecotoxicity, freshwater - inorganics	CTUe	36,5	0,342
Ecotoxicity, freshwater - organics part 1	CTUe	2,88	0,0829
Ecotoxicity, freshwater - organics part 2	CTUe	1,66	0,0097
Particulate matter	disease inc.	0,00000215	0,0000000128
Eutrophication, marine	kg N eq	0,00479	0,0000454
Eutrophication, freshwater	kg P eq	0,000236	0,00000414
Eutrophication, terrestrial	mol N eq	0,0505	0,000334
Human toxicity, cancer	CTUh	0,000000129	0,00000000114
Human toxicity, cancer - inorganics	CTUh	0,00000000548	0,000000000359
Human toxicity, cancer - organics	CTUh	0,000000124	0,0000000011
Human toxicity, non-cancer	CTUh	0,000000419	0,00000000324
Human toxicity, non-cancer - inorganics	CTUh	0,000000367	0,00000000296
Human toxicity, non-cancer - organics	CTUh	0,0000000517	0,000000000281
Ionising radiation	kBq U ⁻²³⁵ eq	0,523	0,00184
Land use	Pt	22,1	0,184
Ozone depletion	kg CFC11 eq	0,00000743	0,000000176
Photochemical ozone formation	kg NMVOC eq	0,0214	0,000131
Resource use, fossils	MJ	81,1	0,557
Resource use, minerals and metals	kg Sb eq	0,0000627	0,00000225
Water use	m ³ depriv.	1,99	0,0105

